arXiv:1503.05238v1 [math.OC] 17 Mar 2015

# Limit value for optimal control with general means*

Xiaoxi LI [†], Marc QUINCAMPOIX [‡], Jérôme RENAULT [§]

March 19, 2015

## Abstract

We consider optimal control problem with an integral cost which is a mean of a given function. As a particular case, the cost concerned is the Cesàro average. The limit of the value with Cesàro mean when the horizon tends to infinity is widely studied in the literature. We address the more general question of the existence of a limit when the averaging parameter converges, for values defined with means of general types.

We consider a given function and a family of costs defined as the mean of the function with respect to a family of probability measures – the evaluations – on $\mathbb{R}_+$. We provide conditions on the evaluations in order to obtain the uniform convergence of the associated value function (when the parameter of the family converges).

Our main result gives a necessary and sufficient condition in term of the total variation of the family of probability measures on $\mathbb{R}_+$. As a byproduct, we obtain the existence of a limit value (for general means) for control systems having a compact invariant set and satisfying suitable nonexpansive property.

**Key words.** optimal control, limit value, general means, long time average value

**AMS subject classifications.** 49J15, 93C15, 37A99

---

[*] This research was partially supported by Agence National de la Researche (grant ANR-10-BLAN 0112). This article is done as part of the PhD thesis of the first author, he wishes to thank his supervisor Sylvain Sorin for advising and comments.

[†] CNRS, IMJ-PRG, UMR 7586, Sorbonne Universités, UPMC Univ Paris 06, Univ Paris Diderot, Sorbonne Paris Cité, Case 247, 4 Place Jussieu, 75252 Paris, France.

[‡] Laboratoire de Mathématiques de Bretagne Atalantique, UMR 6205, Université de Brest, 6 Avenue Le Gorgeu, 29200 Brest, France.

[§] TSE (GREMAQ, Université Toulouse 1 Capitole), 21 allée de Brienne, 31000 Toulouse, France.

# 1 Introduction

We consider a control system defined on $\mathbb{R}^d$ whose dynamic is given by

$$y'(t) = f\big(y(t), u(t)\big) \tag{1.1}$$

where $f : \mathbb{R}^d \times U \to \mathbb{R}^d$ and $u(\cdot)$ is a measurable function – called the control – from $\mathbb{R}_+$ to $U$ a fixed metric space. We will make later on assumptions on (1.1) ensuring that for any initial condition $y(0) = y_0$, and any measurable control $u(\cdot)$, the equation (1.1) has a unique solution $t \mapsto y(t, u, y_0)$ defined on $\mathbb{R}_+$.

To any pair $\big(y_0, u(\cdot)\big)$, we associate a cost

$$\int_0^{+\infty} g\big(y(t, u, y_0), u(t)\big)\mathrm{d}\theta(t),$$

where $g : \mathbb{R}^d \times U \to \mathbb{R}$ is Borel measurable bounded and $\theta \in \Delta(\mathbb{R}_+)$ is a Borel probability measure on $\mathbb{R}_+$. We called $\theta$ an *evaluation* throughout the article.

We will refer to the previously described optimal control problem by the short notation $\mathcal{J} = \langle U, g, f\rangle$. Let $\theta \in \Delta(\mathbb{R}_+)$, we define for $\mathcal{J} = \langle U, g, f\rangle$ the following value function:

$$V_\theta(y_0) = \inf_{u(\cdot)\in\mathcal{U}} \int_0^{+\infty} g\big(y(t, u, y_0), u(t)\big)\mathrm{d}\theta(t), \tag{1.2}$$

where $\mathcal{U}$ denotes the set of measurable controls $u : [0, +\infty) \to U$.

Typical means in the definition (1.2) of the value function are well studied in the literature for

*Cesàro mean*: $\forall t > 0$, $\overline{\theta}_t$ with density $s \mapsto f_{\overline{\theta}_t}(s) = \frac{1}{t}\mathbb{1}_{[0,t]}(s)$, and the $t$-horizon value is

$$V_{\overline{\theta}_t}(y_0) = \inf_{u(\cdot)\in\mathcal{U}} \frac{1}{t}\int_0^t g\big(y(s, u, y_0), u(s)\big)\mathrm{d}s$$

*Abel mean*: $\forall \lambda \in (0, 1]$, $\theta_\lambda$ with density $s \mapsto f_{\theta_\lambda}(s) = \lambda e^{-\lambda s}$, and the $\lambda$-discounted value is

$$V_{\theta_\lambda}(y_0) = \inf_{u(\cdot)\in\mathcal{U}} \int_0^{+\infty} \lambda e^{-\lambda s} g\big(y(s, u, y_0), u(s)\big)\mathrm{d}s$$

The limit of the above value functions as $t$ tends to infinity or as $\lambda$ tends to zero are well investigated in the control literature, (cf. [1], [2], [3], [4], [6], [7] and the references therein), which are often called ergodic control.

When $\theta \in \Delta(\mathbb{R}_+)$ is given, the contribution of the interval $[T, +\infty)$ in the mean (1.2) is less and less significant as $T$ becomes large. Thus the control problem is essentially interesting only on $[0, T_0]$ for certain $T_0$, which we roughly name the "duration" for the problem. In this article, we are interested in the long-run property of $\mathcal{J}$, *i.e.*, the asymptotic behavior of the function $\theta \mapsto V_\theta$ when the "duration" of $\theta$ tends to infinity. In the particular examples of Cesàro mean and Abel mean, the uniform convergence of $V_{\overline{\theta}_t}$ as $t$ tends to infinity and of $V_{\theta_\lambda}$ as $\lambda$ tends to 0 are studied. It is a priori unclear how to define the "duration" of a general evaluation $\theta$ over $\mathbb{R}_+$. If one just assumes the expectation of $\theta$ to be large, we can obtain very different value functions, as is shown by the following

**Example 1.1** *Consider the uncontrolled dynamic* $y(t) = t$, *the running cost* $t \mapsto g(t) = \mathbb{1}_{\cup_{m=1}^{\infty}[2m-1,2m]}(t)$, *and two sequences of evaluations* $(\mu^k)_{k\geq 1}$ *and* $(\nu^k)_{k\geq 1}$ *with densities:* $f_{\mu^k} = \frac{1}{k}\mathbb{1}_{\cup_{m=1}^{k}[2m-1,2m]}$ *and* $f_{\nu^k} = \frac{1}{k}\mathbb{1}_{\cup_{m=1}^{k}[2m-2,2m-1]}$. *Clearly,* $V_{\mu^k} = 1$ *and* $V_{\nu^k} = 0$, $\forall k \geq 1$.

For this reason, we introduce an asymptotic regularity condition for evaluations, called the *long-term condition* (LTC for short), to express the "large duration" and the "asymptotic uniformity of distributions over $\mathbb{R}_+$", and we will study the convergence of the value functions along a sequence of evaluations satisfying the LTC.

More precisely, for any $s \geq 0$, we define the *$s$-total variation* of an evaluation $\theta$ to be the total variation between the measure $\theta$ and its $s$-shift along $\mathbb{R}_+$:

$$TV_s(\theta) = \max_{Q \in \mathcal{B}(\mathbb{R}_+)} |\theta(Q) - \theta(Q+s)|.$$

We say that a sequence of evaluations $(\theta^k)_{k\geq 1}$ satisfies the LTC if:

$$\forall S > 0, \ \sup_{0 \leq s \leq S} TV_s(\theta^k) \xrightarrow[k\to\infty]{} 0.$$

The optimal control problem $\mathcal{J} = \langle U, g, f \rangle$ has a *general limit value* given by some function $V^*$ defined on $\mathbb{R}^d$ if for any sequence $(\theta^k)_k$ satisfying the LTC, $(V_{\theta^k}(y_0))_k$ converges uniformly to $V^*$ as $k$ tends to infinity.

Our main result (Theorem 4.1) states that for any $(\theta^k)_k$ satisfying the LTC, $(V_{\theta^k})_k$ converges uniformly if and only if the family $\{V_{\theta^k}\}$ is totally bounded with respect to the uniform norm. Moreover, in this case, the limit is characterized by the following:

$$V^*(y_0) =_{def} \sup_{\theta \in \Delta(\mathbb{R}_+)} \inf_{s \in \mathbb{R}_+} \inf_{u \in \mathcal{U}} \int_0^{\infty} g\Big(y(t+s, u, y_0), u(t+s)\Big) \mathrm{d}\theta(t), \quad \forall y_0 \in \mathbb{R}^d. \tag{1.3}$$

The above function $V^*$ naturally appears to be the unique possible long-term value function of the control problem.

As a byproduct of our main result, we obtain the existence of the general limit value for any control problem $\mathcal{J} = \langle U, g, f \rangle$ with a running cost $g$ that does not depend on $u$ and with a control dynamic (1.1) which is non-expansive and has a compact invariant set. This can be viewed as a generalization of already obtained results in [8] for optimal control with Cesàro mean.

Existing results in the erdogic control literature are concerned mainly with the convergence of the $t$-horizon Cesàro mean values or the convergence of the $\lambda$-discounted Abel mean values. To the best of the authors' knowledge, this paper is the first to consider general long-term evaluations for optimal control problems.

Also it is worth pointing out that while many works (including [1], [2], [3], [4], [6], [7]) suppose controllability or ergodicity conditions, the present approach does not reply on such conditions. This could be understood by the fact that the limit value $V^*$ may depend on the initial state $y_0$ (which does not occur under ergodic or controlability assumptions).

We also make here a link to the discrete time framework, in which an evaluation $\theta = (\theta_m)_{m\geq 1}$ is a probability measure over positive integers $\mathbb{N}^* = \mathbb{N}\backslash\{0\}$, and $\theta_t$ is the weight for the stage-$t$ payoff. The analogue notion of *total variation* is defined for any $\theta \in \Delta(\mathbb{N}^*)$: $TV(\theta) =$

$\sum_{m=1}^{\infty} |\theta_{m+1} - \theta_m|$ (cf. [12] and [9]). Recently, the existence of the general limit value of dynamic optimization problems in several discrete time frameworks has been obtained in [9], [11] and [13]. Our work is partially inspired by [9] and similar tool within the proof appeared in [10].

The article is organized as follows. Section 2 contains some preliminary notations and basic examples. The long-term condition is introduced and studied in Section 3. Section 4 contains our main result and its consequences. We discuss in the end of this section two (counter)examples. Section 5 is devoted to the proof of the main result. A weaker notation of LTC is discussed in Section 6.

## 2 Preliminaries

Consider the optimal control problem $\mathcal{J} = \langle U, g, f \rangle$ described by (1.1)-(1.2). We make the following assumptions on $g$ and $f$:

$$\begin{cases} \text{the function } g : \mathbb{R}^d \times U \to \mathbb{R} \text{ is Borel measurable and bounded;} \\ \text{the function } f : \mathbb{R}^d \times U \to \mathbb{R}^d \text{ is Borel measurable, and satisfies:} \\ (*).\ \exists L \geq 0, \forall (y, \overline{y}) \in \mathbb{R}^{2d}, \forall u \in U, ||f(y,u) - f(\overline{y}, u)|| \leq L||y - \overline{y}||, \\ (**).\ \exists a > 0, \forall (y,u) \in \mathbb{R}^d \times U, ||f(y,u)|| \leq a(1 + ||y||). \end{cases} \tag{2.1}$$

Under these hypotheses, given any control $u(\cdot)$ in $\mathcal{U}$ and any initial starting state $y_0 \in \mathbb{R}^d$, (1.1) has a unique absolutely continuous solution $t \mapsto y(t, u, y_0)$ defined on $[0, +\infty)$. As the running cost function $g : \mathbb{R}^d \times U \to \mathbb{R}$ is bounded, we can always assume that $g : \mathbb{R}^d \times U \to [0,1]$ after some affine transformation.

Below we introduce several notations.

**$\theta$-evaluated cost $\gamma_\theta(y_0, u)$** Given $\theta \in \Delta(\mathbb{R}_+)$ and $y_0 \in \mathbb{R}^d$, the $\theta$-evaluated cost induced by a control $u(\cdot) \in \mathcal{U}$ is denoted by:

$$\gamma_\theta(y_0, u) = \int_{[0,+\infty)} g\Big(y(s, u, y_0), u(s)\Big) \mathrm{d}\theta(s),$$

With this notation, the $\theta$-value function in (1.2) writes as $V_\theta(y_0) = \inf_{u(\cdot) \in \mathcal{U}} \gamma_\theta(y_0, u)$.

**Reachable map $R_t$** For any $y_0 \in \mathbb{R}^d$, the reachable map in $\mathbb{R}_+$, $t \mapsto R_t(y_0)$, is defined as:

$$R_t(y_0) = \Big\{ \overline{y} \in \mathbb{R}^d \big| \exists\ u(\cdot) \in \mathcal{U} : y(t, u, y_0) = \overline{y} \Big\}. \tag{2.2}$$

$R_t(y_0)$ represents the set of states that the dynamic can reach via certain control at time $t$, starting from the initial state $y_0$ at time 0. We write $R^t(y_0) = \cup_{s=0}^{t} R_s(y_0)$ and $R(y_0) = \cup_{s=0}^{\infty} R_s(y_0)$. $R(y_0)$ is the set of states that can be reached at any finite time starting from $y_0$.

**Image measure $\mathcal{T}_t \sharp \theta$ and the auxiliary value function $V_{\mathcal{T}_t \sharp \theta}$** Given $t \in \mathbb{R}$ and $\theta$ in $\Delta(\mathbb{R}_+)$, we use $\mathcal{T}_t \sharp \theta$ to denote the image (push-forward) measure of $\theta$ by the function $\mathcal{T}_t : s \mapsto s + t$, *i.e.*,

$$\mathcal{T}_t \sharp \theta(Q) = \theta\left(\mathcal{T}_t^{-1}(Q)\right) = \theta\big((Q - t) \cap \mathbb{R}_+\big),\ \forall Q \in \mathcal{B}(\mathbb{R}_+),$$

where $\mathcal{B}(\mathbb{R}_+)$ denotes the set of all Borel subsets in $\mathbb{R}_+$. This leads us to write the $t$-shift $\theta$-evaluated cost induced by a control $u$ as follow:

$$\gamma_{\mathcal{T}_t\sharp\theta}(y_0, u) = \int_{[0,+\infty)} g\Big(y(s+t, u, y_0), u(s+t)\Big)\mathrm{d}\theta(s), \ \forall t \geq 0. \tag{2.3}$$

Taking on both sides of (2.3) the infimum over $u(\cdot) \in \mathcal{U}$ and using the notation of reachable map $R_t$, we obtain the $t$-shift $\theta$-value function

$$V_{\mathcal{T}_t\sharp\theta}(y_0) = \inf_{u(\cdot)\in\mathcal{U}} \int_{[0,+\infty)} g\Big(y(s+t, u, y_0), u(s+t)\Big)\mathrm{d}\theta(s) = \inf_{\overline{y}\in R_t(y_0)} V_\theta(\overline{y}). \tag{2.4}$$

**s-total variation** Given an evaluation $\theta$, define its $s$-total variation for each $s \geq 0$:

$$TV_s(\theta) = \sup_{Q\in\mathcal{B}(\mathbb{R}_+)} |\theta(Q) - \theta(Q+s)|. \tag{2.5}$$

**Long-term condition (LTC)** A sequence of evaluations $(\theta^k)_{k\geq 1}$ satisfies the LTC if:

$$\forall S > 0, \ \overline{TV}_S(\theta^k) =_{def} \sup_{0\leq s\leq S} TV_s(\theta^k) \xrightarrow[k\to\infty]{} 0. \tag{2.6}$$

In this article, we are concerned with the following notation of limit value for optimal control problems with general means.

**Definition 2.1** *Let $V$ be a function defined on $\mathbb{R}^d$. The optimal control problem $\mathcal{J}$ admits $V$ as the **general limit value** if: for any sequence of evaluations $(\theta^k)_{k\geq 1}$ satisfying the LTC, for all $y_0$ in $\mathbb{R}^d$, $(V_{\theta^k}(y_0))$ converges to $V(y_0)$ as $k$ tends to infinity, and moreover the convergence is uniform in $y_0$.*

Below are some basic examples of optimal control problems in which the general limit value exists.

**Example 2.2** *$y$ lies in $\mathbb{R}^2$ seen as the complex plane, there is no control, and the dynamic is given by $f(y) = i\ y$, where $i^2 = -1$. We clearly have*

$$V_{\theta^k}(y_0) \xrightarrow[k\to\infty]{} \frac{1}{2\pi}\int_0^{2\pi} g(|y_0|e^{rit})\mathrm{d}t,$$

*for any sequence of evaluations $(\theta^k)_k$ satisfying the LTC.*

**Example 2.3** *$y$ lies in the complex plane again, with $f(y,u) = i\ y\ u$, where $u \in U$ is a given bounded subset of $\mathbb{R}$, and $g$ is any continuous function in $y$ (which thus does not depend on $u$).*

**Example 2.4** *$f(y,u) = -y + u$, where $u \in U$ a given bounded subset of $\mathbb{R}^d$, and $g$ is any continuous function in $y$ (which thus does not depend on $u$).*

We will show later (using Corollary 4.7) that the general limit value exists in Examples 2.3 and 2.4.

# 3 On the long-term condition (LTC)

In this section, we discuss the LTC. First, we give the following remarks.

**Remark 3.1** *(a). By definition, one has*

$$\forall s \geq 0, \forall t \geq 0, \forall \theta \in \Delta(\mathbb{R}_+),\ TV_{s+t}(\theta) \leq TV_s(\theta) + TV_t(\theta).$$

*This implies that $(\theta^k)_{k\geq 1}$ satisfies the LTC if and only if $\overline{TV}_1(\theta^k) \xrightarrow[k\to\infty]{} 0$.*
*(b). If one takes $Q = \mathbb{R}_+$ in definition of $TV_s(\theta^k)$ for each $s \geq 0$ and each $k \geq 1$, we deduce that if $(\theta^k)_{k\geq 1}$ satisfies the LTC, then $\theta^k([0,s]) \xrightarrow[k\to\infty]{} 0$ for any $s \geq 0$.*

**Remark 3.2** *Let $\theta$ be an evaluation absolutely continuous w.r.t. the Lebesgue measure on $\mathbb{R}_+$, and $f_\theta$ its density. Scheffé Theorem (cf. [5], Theorem 1 in p.2) implies that:*

$$\forall s \geq 0,\ 2TV_s(\theta) = I_s(\theta) =_{def} \int_0^\infty |f_\theta(t+s) - f_\theta(t)| \mathrm{d}t.$$

*Thus, if $(\theta^k)_{k\geq 1}$ is a sequence of evaluations with densities $(f_{\theta^k})_{k\geq 1}$:*
*(a). $(\theta^k)_{k\geq 1}$ satisfies the LTC if and only if $\sup_{0\leq s\leq 1} I_s(\theta^k) \xrightarrow[k\to\infty]{} 0$. If moreover, for each $k \geq 1$, $t \mapsto f_{\theta^k}(t)$ is non increasing on $\mathbb{R}_+$, then $(\theta^k)_{k\geq 1}$ satisfies the LTC if and only if $\forall s \geq 0$, $\theta^k([0,s]) = \int_{t=0}^\infty f_\theta(t)\mathrm{d}t - \int_{t=0}^\infty f_\theta(t+s)\mathrm{d}t \xrightarrow[k\to\infty]{} 0$.*
*(b). if $(\theta^k)_{k\geq 1}$ satisfies the LTC, then $\int_{t=0}^\infty t f_{\theta^k}(t)\mathrm{d}t \xrightarrow[k\to\infty]{} \infty$. Indeed, Chebychev's inequality gives that $\int_{t=0}^\infty t f_{\theta^k}(t)\mathrm{d}t \geq M\left(1 - \theta^k([0,M])\right)$ for all $M > 0$.*

Here we discuss several cases where the LTC condition is satisfied.

**Example 3.3** *(Uniform distributions) Assume that for each $k$, $\theta^k$ is the uniform law over the interval $[a_k, b_k]$, with $0 \leq a_k \leq b_k$. For each $k$,*

- $\underline{s \geq b_k - a_k}$: $I_s(\theta^k) = \begin{cases} \frac{2}{b_k - a_k} & \text{if} \quad 0 < s < a_k \\ \frac{1+(b_k - s)}{b_k - a_k} & \text{if} \quad a_k < s < b_k \\ \frac{1}{b_k - a_k} & \text{if} \quad b_k < s \end{cases}$,
- $\underline{s < b_k - a_k}$: $I_s(\theta^k) = \begin{cases} \frac{2s}{b_k - a_k} & \text{if} \quad 0 < s < a_k \\ \frac{2s}{b_k - a_k} & \text{if} \quad a_k < s < b_k \end{cases}$.

*One can check easily that $(\theta^k)_k$ satisfies the LTC if and only if $b_k - a_k \xrightarrow[k\to\infty]{} \infty$. Indeed, by Remark 3.2 (a), it is sufficient to look at $I_s(\theta^k)$ for $s \in [0,1]$.*

**Example 3.4** *(Abel average) Assume that for each $k$, $\theta^k$ has density $s \mapsto f_{\theta^k}(s) = \lambda_k e^{-\lambda_k s}\mathbb{1}_{\mathbb{R}_+}(s)$, with $\lambda_k > 0$. Since $\forall k \geq 1$, $s \mapsto f_{\theta^k}(s)$ is non increasing, Remark 3.2 (a) implies that $(\theta^k)_k$ satisfies the LTC if and only if: $\forall T > 0$, $\theta^k([0,T]) = \int_{s=0}^T \lambda_k e^{-\lambda_k s}\mathrm{d}s = 1 - e^{-T/\lambda_k} \xrightarrow[k\to\infty]{} 0$, which is again equivalent to $\lambda_k \xrightarrow[k\to\infty]{} 0$.*

**Example 3.5** *(Folded normal distributions) Assume that for each $k$, $\theta^k$ is the distribution of a random variable $|X^k|$, where $X^k$ follows a normal law $\mathcal{N}(m_k, \sigma_k^2)$. The density of $\theta^k$ is given by:*

$$\forall t \geq 0,\ f_{\theta^k}(t) = \frac{1}{\sigma_k\sqrt{2\pi}}\left[\exp\left(-\frac{1}{2}\Big(\frac{t-m_k}{\sigma_k}\Big)^2\right) + \exp\left(-\frac{1}{2}\Big(\frac{t+m_k}{\sigma_k}\Big)^2\right)\right].$$

**Claim 3.1** *$(\theta^k)_k$ satisfies the LTC if and only if $\sigma_k \xrightarrow[k\to\infty]{} \infty$.*

Our argument relies on the following lemma, whose proof is put in the **Appendix**. Without loss of generality, we may assume that $m_k$ is non-negative for each $k$.

**Lemma 1** *Let $\theta$ be the distribution of $X$ where $|X|$ follows the normal law $\mathcal{N}(m, \sigma)$ with $m, \sigma > 0$. There exists some $t^* \in [0, m)$ such that $f'_\theta(t) > 0$ for any $t \in (0, t^*)$ and $f'_\theta(t) < 0$ for any $t \in (t^*, \infty)$. Moreover, such $t^*$ satisfies that: $(t^*)^2 \geq m^2 - \sigma^2$.*

**Proof of Claim 3.1** We apply Lemma 1 to each evaluation $\theta_k$ to obtain some $t_k^* \in [0, m_k)$ such that: $f_{\theta^k}(\cdot)$ is increasing on $[0, t_k^*)$ and decreasing on $[t_k^*, \infty)$. This enables us to write:

$$\forall s \leq t_k^*,\ \ I_s(\theta^k) = \int_{t_k^*-s}^{t_k^*} f_{\theta^k}(t)\mathrm{d}t + \int_{t_k^*-s}^{t_k^*} |f_{\theta^k}(t+s) - f_{\theta^k}(t)|\mathrm{d}t + \int_{t_k^*}^{t_k^*+s} f_{\theta^k}(t)\mathrm{d}t.$$

We deduce then $sf_{\theta^k}(t_k^* - s) \leq I_{\theta^k}(s) \leq 4sf_{\theta^k}(t_*^k)$ for $s \leq t_k^*$. Assume below $\hat{t}^* =_{def} \liminf_{k\to\infty} t_k^* > 0$, and the analysis is analogue for $\hat{t}^* = 0$, which we omit here.
(*). Suppose that $\sigma_k \to \infty$, then

$$f_{\theta^k}(t_k^*) = \frac{1}{\sigma_k\sqrt{2\pi}}\left[\exp\left(-\frac{1}{2}\Big(\frac{t_k^*-m_k}{\sigma_k}\Big)^2\right) + \exp\left(-\frac{1}{2}\Big(\frac{t_k^*+m_k}{\sigma_k}\Big)^2\right)\right] \leq \frac{2}{\sigma_k\sqrt{2\pi}} \xrightarrow[k\to\infty]{} 0.$$

This implies that for $S = \hat{t}^* \wedge 1$, $\sup_{0\leq s\leq S} I_s(\theta^k) \xrightarrow[k\to\infty]{} 0$.
(**). Conversely, suppose that $(\theta^k)_k$ satisfies the LTC. Then for any $s < \hat{t}^*$, $I_s(\theta^k)$ thus $f_{\theta^k}(t_k^* - s)$ vanishes as $k$ tends to infinity. This implies that either $\sigma_k \to \infty$ or $(\sigma_k)_k$ is bounded and $\big(m_k - (t_k^* - s)\big)_k \to \infty$. Lemma 1 shows that the specified point $t_k^*$ for the evaluation $\theta_k$ satisfies $(t_k^*)^2 \geq m_k^2 - \sigma_k^2$, thus $m_k - (t_k^* + s) \leq m_k - t_k^* \leq \frac{\sigma_k^2}{m_k + t_k^*} \leq \frac{\sigma_k^2}{m_k}$. If $(\sigma_k)_k$ is bounded, $(m_k - t_k^*)_k$ thus $(m_k)_k$ should tend to infinity, but this leads to a contradiction with $m_k - t_k^* \leq \frac{\sigma_k^2}{m_k}$. □

Now we link the LTC condition to the discrete time framework. In a discrete time dynamic optimization problem, a general evaluation on the payoff stream is a probability distribution over $\mathbb{N}^* = \mathbb{N}/\{0\}$ the set of postive integers. For any $\xi = (\xi_1, ..., \xi_t, ...)$ in $\Delta(\mathbb{N}^*)$, its "total variation" $TV(\xi) = \sum_{m=1}^\infty |\xi_{m+1} - \xi_m|$ is the stage by stage absolute difference between the measure $\xi$ and its one-stage "shift" measure $\xi' = (\xi_2, ..., \xi_{t+1}, ...)$. (cf. Sorin [12] or Renault [11]).

When the sequence of evaluations in continuous time admits step functions as densities, this link to discrete time framework is much clearer as seen by the following

**Proposition 3.6** *Let $(\theta^k)_k$ be a sequence of absolutely continuous evaluations in $\Delta(\mathbb{R}_+)$, and their densities are given as: $\forall k \geq 1, f_{\theta^k} = \sum_{m=1}^\infty \xi_m^k \mathbb{1}_{[m-1,m)}$, where $\xi^k = (\xi_1^k, ..., \xi_m^k, ...,) \in \Delta(\mathbb{N}^*)$. Then $(\theta^k)_k$ satisfies the LTC if and only if $\sum_{m=1}^\infty |\xi_{m+1}^k - \xi_m^k| \xrightarrow[k\to\infty]{} 0$.*

**Proof**: Fix $s \in [0,1]$. We shall write for each $k$,

$$I_s(\theta^k) = \sum_{m=1}^{\infty} \int_{[m-1,m)} \Big| f_{\theta^k}(t+s) - f_{\theta^k}(t) \Big| \mathrm{d}t.$$

For each $m = 1, 2, ...$, we have

$$\begin{aligned}\int_{[m-1,m)} \Big| f_{\theta^k}(t+s) - f_{\theta^k}(t) \Big| \mathrm{d}t =& \int_{[m-1,m-s)} \Big| f_{\theta^k}(t+s) - f_{\theta^k}(t) \Big| \mathrm{d}t + \int_{[m-s,m)} \Big( f_{\theta^k}(t+s) - f_{\theta^k}(t) \Big) \mathrm{d}t \\ =& s\big|\xi^k_{m+1} - \xi^k_m\big|.\end{aligned}$$

As a consequence,

$$I_s(\theta^k) = s \sum_{m=1}^{\infty} \big|\xi^k_{m+1} - \xi^k_m\big| \leq \sum_{m=1}^{\infty} \big|\xi^k_{m+1} - \xi^k_m\big|,\ \forall s \in [0,1].$$

In view of Remark 3.2, $(\theta^k)_k$ satisfies the LTC if and only if $\sum_{m=1}^{\infty} |\xi^k_{m+1} - \xi^k_m| \xrightarrow[k\to\infty]{} 0$. $\square$

We end this section by a preliminary lemma, which will be useful in later results.

**Lemma 3.7** *Fix any* $\theta \in \Delta(\mathbb{R}_+)$ *and any* $t \in \mathbb{R}_+$, *we have*

$$\left| \int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s-t)\mathrm{d}\theta(s) \right| \leq TV_t(\theta) \textit{ and } \left| \int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s+t)\mathrm{d}\theta(s) \right| \leq 2TV_t(\theta),$$

*for any* $h(\cdot) \in \mathcal{M}(\mathbb{R}_+, [0,1])$, *where* $\mathcal{M}(\mathbb{R}_+, [0,1]) = \Big\{ h(\cdot) \Big| h : \mathbb{R}_+ \to [0,1], \textit{Borel measurable} \Big\}$.

**Proof**: We fix any $\theta \in \Delta(\mathbb{R}_+)$ and $t \in \mathbb{R}_+$. By definition of $\mathcal{T}_s\sharp\theta$, we have that for any $h(\cdot) \in \mathcal{M}(\mathbb{R}_+, [0,1])$:

$$\int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[t,+\infty)} h(s-t)\mathrm{d}\theta(s) = \int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s)\mathrm{d}\mathcal{T}_{-t}\sharp\theta(s) \tag{3.1}$$

and

$$\int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s+t)\mathrm{d}\theta(s) = \int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s)\mathrm{d}\mathcal{T}_{t}\sharp\theta(s). \tag{3.2}$$

Since $\mathcal{T}_{-t}\sharp\theta$ and $\mathcal{T}_t\sharp\theta$ are both Borel measures on $\mathbb{R}_+$, "$\theta - \mathcal{T}_{-t}\sharp\theta$" and "$\theta - \mathcal{T}_t\sharp\theta$" are both signed measures. Hahn's decomposition theorem[1] implies that:

$$\sup_{h \in \mathcal{M}(\mathbb{R}_+,[0,1])} \left| \int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s)\mathrm{d}\mathcal{T}_{-t}\sharp\theta(s) \right| = \sup_{Q \in \mathcal{B}(\mathbb{R}_+)} \Big| \theta(Q) - \mathcal{T}_{-t}\sharp\theta(Q) \Big|.$$

and

$$\sup_{h \in \mathcal{M}(\mathbb{R}_+,[0,1])} \left| \int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s)\mathrm{d}\mathcal{T}_{t}\sharp\theta(s) \right| = \sup_{Q \in \mathcal{B}(\mathbb{R}_+)} \Big| \theta(Q) - \mathcal{T}_{t}\sharp\theta(Q) \Big|.$$

---

[1] The first author acknowledges Eilon Solan for the discussion on using Hahn's decomposition theorem.

Combining with (3.1)-(3.2), we obtain:

$$\left|\int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[t,+\infty)} h(s-t)\mathrm{d}\theta(s)\right| \leq \sup_{Q\in\mathcal{B}(\mathbb{R}_+)} \Big|\theta(Q)-\theta(Q+t)\Big| = TV_t(\theta)$$

and

$$\left|\int_{[0,+\infty)} h(s)\mathrm{d}\theta(s) - \int_{[0,+\infty)} h(s+t)\mathrm{d}\theta(s)\right| \leq \sup_{Q\in\mathcal{B}(\mathbb{R}_+)} \Big|\theta(Q)-\theta(Q-t)\Big| \leq \theta\big([0,t)\big) + TV_t(\theta) \leq 2TV_t(\theta).$$

The proof of the lemma is complete. □

## 4 Main Result

As will be shown in our main result, the function $V^*(y_0)$ defined in (1.3) characterizes the general limit value of the optimal control problem in case of convergence. We first rewrite it as

$$V^*(y_0) = \sup_{\mu\in\Delta(\mathbb{R}_+)} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\mu}(y_0) = \sup_{\mu\in\Delta(\mathbb{R}_+)} \inf_{\overline{y}\in R(y_0)} V_\mu(\overline{y}).$$

We give the following interpretation: consider the auxiliary optimal control problem (game) where an adversary of the controller chooses an evaluation $\mu$, and then knowing $\mu$ as given, the controller chooses an initial state in the reachable set $R(y_0)$. The running cost from time $t$ is evaluated by $\mu$ and $V^*(y_0)$ is the value of this problem starting from $y_0$.

Recall that a metric space $X$ is *totally bounded* if for each $\varepsilon > 0$, $X$ can be covered by finitely many balls of radius $\varepsilon$.

**Theorem 4.1** *Let $(\theta^k)_{k\geq 1}$ be a sequence of evaluations satisfying the LTC. Assume (2.1) for the optimal control problem $\mathcal{J} = \langle U, g, f\rangle$. Then,*
*(i). $V^* = \sup_{k\in\mathbb{N}} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^k}$.*
*(ii). Any accumulation point (for the uniform convergence) of the sequence $(V_{\theta^k})_k$ is equal to $V^*$.*
*(iii). The sequence $(V_{\theta^k})_k$ uniformly converges if and only if the space $(\{V_{\theta^k}\}, ||\cdot||_\infty)$ is totally bounded.*

**Remark 4.2** *Let $(\theta^k)_k$ be a sequence of evaluations which contains a subsequence $(\theta^{\varphi_k})_k$ satisfying the LTC. Then Part (i) of Theorem 4.1 still holds true for $(\theta^k)_k$.*

A more precise convergence result is obtained if we suppose that there exists a compact set $Y \subseteq \mathbb{R}^d$ which is *invariant* for the dynamic (1.1), *i.e.*, such that $y(t, u, y_0) \in Y$ for all $u(\cdot) \in \mathcal{U}$, $t \geq 0$ and $y_0$ in $Y$.

**Definition 4.3** *Let $V$ be a function defined on $\mathbb{R}^d$. In the optimal control problem $\mathcal{J}$, there is the **general uniform convergence** of the value functions $\{V_\theta\}$ to $V$ if:*

$$\forall\varepsilon > 0, \exists S > 0, \exists\eta > 0 \text{ s.t. } \forall\theta\in\Delta(\mathbb{R}_+), \text{with } \overline{TV}_S(\theta) \leq \eta,\ ||V_\theta - V||_\infty \leq \varepsilon.$$

**Lemma 4.4** *Let $V$ be a function defined on $\mathbb{R}^d$. The optimal control problem $\mathcal{J}$ admits $V$ as the general limit value if and only if there is the general uniform convergence of the value functions $\{V_\theta\}$ to $V$.*

**Proof.** The general uniform convergence of the value functions $\{V_\theta\}$ to $V$ implies the existence of general limit value given as $V$. Next we show that the existence of the general limit value given as $V$ is sufficient to deduce the general uniform convergence of $\{V_\theta\}$ to $V$. Suppose by contradiction that there is no general uniform convergence of $\{V_\theta\}$ to $V$, *i.e.*,

$$\exists \varepsilon_0 > 0,\ \forall S > 0,\ \forall \eta^k > 0,\ \exists \theta^k \in \Delta(\mathbb{R}_+) \text{ with } \overline{TV}_S(\theta^k) \le \eta^k, \text{ and } ||V_{\theta^k} - V||_\infty > \varepsilon_0,\ \forall k \ge 1.$$

Let $\varepsilon_0 > 0$ be fixed as above. We take a vanishing positive sequence $(\eta^k)_k$ and some $S_0 > 0$, then there is a sequence of evaluations $(\theta^k)$ with $\overline{TV}_{S_0}(\theta^k) \le \eta^k \xrightarrow[k\to\infty]{} 0$, and $\liminf_k ||V_{\theta^k} - V||_\infty \ge \varepsilon_0$. According to Remark 3.1 (a), such $(\theta^k)_k$ satisfies the LTC, while $(V_{\theta^k})$ does not converges uniformly to $V^*$. This is a contradiction. ■

**Corollary 4.5** *Assume (2.1) for the optimal control problem $\mathcal{J} = \langle U, g, f\rangle$. Suppose that there is a compact set $Y \subseteq \mathbb{R}^d$ which is invariant for the dynamic (1.1), and that the family $\{V_\theta : \theta \in \Delta(\mathbb{R}_+)\}$ is uniformly equicontinuous on $Y$. Then there is the general uniform convergence of the value functions $\{V_\theta\}$ to $V^*$.*

**Proof**: By assumption, the family of value functions $\{V_\theta : \theta \in \Delta(\mathbb{R}_+)\}$ is both uniformly bounded and uniformly equicontinuous on the compact invariant set $Y$, so we can use Ascoli's theorem to deduce the totally boundedness of the space $(\{V_\theta\}, ||\cdot||_\infty)$. Theorem 4.1 implies that: for any $(\theta^k)_k$ satisfying the LTC, the corresponding sequence of value functions $(V_{\theta^k})$ converges uniformly to $V^*$ as $k$ tends to infinity. Thus $\mathcal{J}$ has a general limit value given as $V^*$, and according to Lemma 4.4, there is the uniform convergence of value functions $\{V_\theta\}$ to $V^*$. □

We shall give the existence result of the general limit value under sufficient conditions expressed directly in terms of properties of the control dynamic (1.1) and of the running cost $g$.

Let us introduce the following *non expansive* condition (cf. [8]). The control dynamic (1.1) is non expansive if

$$\forall y_1, y_2 \in \mathbb{R}^d,\ \sup_{a\in U} \inf_{b\in U} \Big\langle y_1 - y_2, f(y_1, a) - f(y_2, b) \Big\rangle \le 0.$$

**Definition 4.6** *The optimal control problem $\mathcal{J} = \langle U, g, f\rangle$ is called* ***compact non expansive*** *if it satisfies the following three conditions:*
$(A.1)$ *there is a compact set $Y \subseteq \mathbb{R}^d$ is the invariant for the dynamic (1.1);*
$(A.2)$ *the running cost function $g(\cdot)$ does not depend on $u \in U$, and is continuous in $y \in \mathbb{R}^d$;*
$(A.3)$ *the control dynamic (1.1) is non expansive on $Y$.*

**Corollary 4.7** *Assume (2.1) for the optimal control problem $\mathcal{J} = \langle U, g, f\rangle$. Suppose that that $\mathcal{J}$ is compact non expansive, then the general limit value exists in $\mathcal{J}$ and is given as $V^*$.*

**Proof**: Under $(A.1)$ and $(A.3)$, Proposition 3.7 in [8] implies that:

$$\forall (y_1, y_2) \in Y^2, \forall u(\cdot) \in \mathcal{U}, \exists v(\cdot) \in \mathcal{U}, s.t.\ \forall t \geq 0,\ ||y(t, u, y_1) - y(t, v, y_2)|| \leq ||y_1 - y_2||. \tag{4.1}$$

We claim that the family $(V_\theta)_{\theta \in \Delta(\mathbb{R}_+)}$ is uniformly equicontinuous on $Y$, thus Corollary 4.5 and Lemma 4.4 apply. Fix any $(y_1, y_2) \in Y^2$, $\theta \in \Delta(\mathbb{R}_+)$, and $\varepsilon > 0$. Let $u$ be $\varepsilon$-optimal for $V_\theta(y_1)$:

$$V_\theta(y_1) \geq \int_{[0,+\infty)} g\Big(y(s, u, y_1)\Big) \mathrm{d}\theta(s) - \varepsilon.$$

According to the non expansive property, there exists $v(\cdot)$ in $\mathcal{U}$ as in (4.1) such that

$$||y(s, u, y_1) - y(s, v, y_2)|| \leq ||y_1 - y_2||,\ \forall s \geq 0. \tag{4.2}$$

By definition, $V_\theta(y_2) \leq \int_{[0,+\infty)} g\big(y(s, y_2, v)\big) \mathrm{d}\theta(s)$, hence

$$V_\theta(y_2) - V_\theta(y_1) \leq \int_{[0,+\infty)} \Big[g\big(y(s, v, y_2)\big) - g\big(y(s, u, y_1)\big)\Big] \mathrm{d}\theta(s) + \varepsilon.$$

Denoting $\omega_g$ the modulus of continuity of $g$, we obtain in view of (4.2):

$$V_\theta(y_2) - V_\theta(y_1) \ \leq \ \int_{[0,+\infty)} \Big[g\big(y(s, v, y_2)\big) - g\big(y(s, u, y_1)\big)\Big] \mathrm{d}\theta(s) + \varepsilon \leq \omega_g(||y_1 - y_2||) + \varepsilon.$$

Interchanging $y_1$ and $y_2$ and taking into account of $\varepsilon > 0$ being arbitrary, we deduce that $(V_\theta)_{\theta \in \Delta(\mathbb{R}_+)}$ is uniformly equicontinuous on the invariant set $Y$. This finishes the proof. $\square$

**Remark 4.8** *Both Example 2.3 and Example 2.4 satisfy conditions of Corollary 4.7, so there is general uniform convergence of the value functions* $\{V_\theta\}$ *(the existence of the general limit value).*

**Remark 4.9** *Our result generalizes Proposition 3.3 in [8] which proved the uniform convergence of the $t$-horizon values in compact non expansive optimal control problems.*

We end this section by presenting two (counter)examples, showing that the results in Theorem 4.1 do not hold if some of their conditions is not satisfied.

The first example is an uncontrolled dynamic. We show that if $(\theta^k)_k$ contains no subsequence satisfying the LTC, then the result in Part (i) of Theorem 4.1 does not hold, *i.e.*, $\sup_{k \geq 1} \inf_{t \geq 0} V_{\mathcal{T}_t \sharp \theta^k}(y_0) < \sup_{\theta \in \Delta(\mathbb{R}_+)} \inf_{t \geq 0} V_{\mathcal{T}_t \sharp \theta}(y_0)$ for some $y_0$ (cf. Remark 4.2).

**Counter-example 4.10** *Consider the uncontrolled dynamic on* $\mathbb{R}$*:* $y(0) = y_0$ *and* $y'(t) = -\left(y(t) - 1\right), \forall t \geq 0$*. The trajectory is then* $y(t) = 1 + (y_0 - 1)e^{-t}$*. The running cost function* $g : \mathbb{R} \to [0, 1]$ *is given by:*

$$g(y) = \begin{cases} 0 & \text{if} \quad y < 0 \\ y & \text{if} \quad 0 \leq y \leq 1 \\ 1 & \text{if} \quad y > 1 \end{cases}$$

*We have that* $V^*(y_0) = \sup_{\theta\in\Delta(\mathbb{R}_+)} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta}(y_0) = 1,\ \forall y_0 \in \mathbb{R}$. *Indeed, let* $y_0$ *be given and fix any* $\varepsilon > 0$*, there is some* $T_\varepsilon > 0$ *such that* $|y(T) - 1| \leq \varepsilon$ *for all* $T \geq T_\varepsilon$. *Take an evaluation* $\theta$ *in* $\Delta(\mathbb{R}_+)$ *with* $\theta([0, T_\varepsilon]) = 0$. *This enables us to deduce that: for all* $t \geq 0$,

$$V_{\mathcal{T}_t\sharp\theta}(y_0) = \int_{[T_\varepsilon,+\infty)} g\big(y(s+t)\big)\mathrm{d}\theta(s) \geq \int_{[T_\varepsilon,+\infty)} g\big(y(T_\varepsilon)\big)\mathrm{d}\theta(s) \geq \big(1 - y(T_\varepsilon)\big)\theta([T_\varepsilon,\infty]) \geq 1 - \varepsilon.$$



***Figure 5.1****: The solution* $y(t) = 1 + (y_0 - 1)e^{-t}$ *to the dynamic is represented by the thick curve. For given* $\varepsilon > 0$*,* $T_\varepsilon > 0$ *is chosen such that* $|y(T_\varepsilon) - 1| = \varepsilon$.

*Consider now any sequence of evaluations* $(\theta^k)_k$ *which does not contain any subsequence satisfying the LTC. Under the assumption that the density* $f_{\theta^k}$ *for each evaluation* $\theta^k$ *is non increasing, we show that Part (i) of Theorem 4.1 is not valid:* $V^* \neq \sup_{k\in\mathbb{N}} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^k}$.

*Indeed, let us take any* $y_0 < 1$ *and suppose that* $\sup_{k\in\mathbb{N}} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^k}(y_0) = V^*(y_0)$*, which is equal to* $1$ *as was proved. Let* $\varphi(k)$ *be a subsequence such that* $\lim_{k\to\infty} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^{\varphi(k)}}(y_0) = 1$. $(\theta^{\varphi(k)})_k$ *does not satisfy the LTC by assumption, so Remark 3.2 (a) implies that there exists some* $T > 0$ *with* $\theta^{\varphi(k)}([0,T]) \nrightarrow 0$. *Let* $\varphi_m$ *be the subsequence of* $\varphi$ *and* $\eta > 0$ *such that* $\theta^{\varphi_m(k)}([0,T]) \xrightarrow[k\to\infty]{} \eta$. *We obtain for any* $k \geq 1$,

$$\begin{aligned} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^{\varphi_m(k)}}(y_0) \leq V_{\theta^{\varphi_m(k)}}(y_0) &= \int_{[0,T]} g\left(y(t)\right)\mathrm{d}\theta^{\varphi_m(k)}(t) + \int_{[T,+\infty]} g\left(y(t)\right)\mathrm{d}\theta^{\varphi_m(k)}(t) \\ &\leq y(T)\theta^{\varphi_k(m)}([0,T]) + \theta^{\varphi_k(m)}([T,\infty]). \end{aligned}$$

*This implies that for such fixed* $y_0 < 1$ *,* $\lim_k \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^{\varphi_m(k)}}(y_0) \leq y(T)\eta + (1 - \eta) < 1$. *This contradicts the assumption that* $\sup_{k\in\mathbb{N}} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^k}(y_0) = 1$*, and our claim is proved.*

In the second example, we study the convergence of the value functions of a control problem along two different sequences of evaluations satisfying the LTC. Along the first sequence, the value functions converge uniformly to $V^*$; while along the second, the value functions pointwisely converge, but not uniformly (thus the family of value functions is not totally bounded for the uniform norm), to a limit function which is different from $V^*$.

**Counter-example 4.11** *Consider the control problem on the state space* $\mathbb{R} = (-\infty, +\infty)$*, where* *the control set is* $U = \{+1, -1\}$*; the dynamic is*[2]*:*

$$f(y,u) = u \text{ for all } (y,u) \in \mathbb{R}_+ \times U \text{ and } f(y,u) = -1 \text{ for all } (y,u) \in \mathbb{R}_-^* \times U,$$

*where* $\mathbb{R}_-^* = \mathbb{R}_-/\{0\}$*; and the running cost function is:*

$$g(y,u) = \begin{cases} +1 & \text{if} \quad u = +1,\ y \geq 0 \\ 0 & \text{if} \quad u = -1,\ y \geq 0 \\ +K & \text{if} \quad \quad y < 0 \end{cases}$$

*Suppose that* $K > 1$ *large enough, so the cost on* $\mathbb{R}_-$ *is positive and high. Whenever the state reaches* $y = 0$*, it is optimal to choose control* $u = +1$ *and this drives the state back to* $\mathbb{R}_+$*; on* $\mathbb{R}_-^*$*, the dynamic is* $f = -1$*, independent of control and state.* $V_\theta(y_0) = K$ *for all* $y_0$ *in* $\mathbb{R}_-^*$ *and* $\theta$ *in* $\Delta(\mathbb{R}_+)$*, so the reduced state space is* $\mathbb{R}_+$*, and we consider value functions defined on it.*

$V^*(y_0) = \sup_\theta \inf_{t \geq 0} V_{\mathcal{T}_t \sharp \theta}(y_0) = 0$ *for any* $y_0 \geq 0$*. Fix any* $y_0 \geq 0$*. For any* $\theta \in \Delta(\mathbb{R}_+)$ *and* $\varepsilon > 0$*, let* $t^\varepsilon \geq 0$ *such that* $\theta([0, t^\varepsilon]) \geq 1 - \varepsilon$*. Define now the control* $u^\varepsilon(\cdot)$ *to be:* $u^\varepsilon(t) = +1$*, if* $t \in [0, t^\varepsilon]$ *and* $u^\varepsilon(t) = -1$ *if* $t \in (t^\varepsilon, \infty)$*, which gives:* $\gamma_{\mathcal{T}_{t^\varepsilon} \sharp \theta}(y_0, u^\varepsilon) \leq \varepsilon K$*.*

*Consider* $(\theta^k)_k$ *the sequence of evaluations with density* $f_{\theta^k}(s) = \frac{1}{k}\mathbb{1}_{[k,2k]}(s)$ *for each* $k$*, and* $(\bar{\theta}^k)_k$ *the sequence of* $k$*-horizon evaluations with density* $f_{\bar{\theta}^k}(s) = \frac{1}{k}\mathbb{1}_{[0,k]}(s)$ *for each* $k$*. We show that:*

$(\{V_{\theta^k}\}, ||\cdot||_\infty)$ *is totally bounded and* $(V_{\theta^k})$ *converges uniformly to* $V^*$*; while* $(\{V_{\overline{\theta}^k}\}, ||\cdot||_\infty)$ *is not totally bounded and* $(V_{\overline{\theta}^k})$ *does not converge to* $V^*$*.*

*Let* $y_0 \geq 0$*, we have that:*

1. $V_{\theta^k}(y_0) = 0$*, for all* $k \geq 1$*. Indeed, one optimal control for* $V_{\theta^k}(y_0)$ *can be taken as:* $u^*(t) = +1,\ t \in [0,k]$ *and* $u^*(t) = -1,\ t \in (k, 2k]$*;*
2. $V_{\overline{\theta}^k}(y_0) = 0$ *if* $k \leq y_0$ *and* $V_{\overline{\theta}^k}(y_0) = \frac{1}{2} - \frac{y_0}{2k}$ *if* $k > y_0$*. Indeed, for* $k \leq y_0$*, one optimal control for* $V_{\overline{\theta}^k}(y_0)$ *can be taken as:* $u^*(t) = -1,\ t \in [0,k]$*; for* $k > y_0$*, one optimal control for* $V_{\overline{\theta}^k}(y_0)$ *can be taken as:* $u^*(t) = +1,\ t \in [0, \frac{k-y_0}{2}]$ *and* $u^*(t) = -1,\ t \in (\frac{k-y_0}{2}, k]$*, so* $\gamma_{\overline{\theta}^k}(y_0, u^*) = \frac{(k-y_0)/2}{k} = \frac{1}{2} - \frac{y_0}{2k}$*.*

*See the following two pictures for illustration.*

---

[2] Notice that the dynamic is discontinuous at $y = 0$ when $u = +1$. To get the desired asymptotic result under the Liptchitz regularity, one can slightly modify dynamic to set $f(y, +1) = y$ for $y \in [0, 1]$ and others unchanged.



***Figure 5.2***: *The left figure describes the dynamic of one optimal control for the evaluation $\theta^k$, which is $u^* = +1$ on $[0, k]$ and $u^* = -1$ on $(k, 2k]$; the right figure describes the dynamic of one optimal control for the evaluation $\bar{\theta}^k$ with $k > y_0$, which is $u^* = +1$ on $[0, \frac{k-y_0}{2}]$ and $u^* = -1$ on $(\frac{k-y_0}{2}, k]$. Here, the vertical axis represents the distance of $y(t)$ from zero and the thick trajectory (resp. thin trajectory) corresponds to state on which $u = +1$ and $g = 1$ (resp. $u = -1$ and $g = 0$).*

*We deduce that $(V_{\theta^k}(y_0))_k$ converges uniformly to $V^*(y_0) = 0$ on $\mathbb{R}_+$; and that $V_{\overline{\theta}^k}(y_0) \xrightarrow[k\to\infty]{} \frac{1}{2}$, while the convergence is not uniformly in $y_0 \in \mathbb{R}_+$: indeed, for all $k \geq 1$, $V_{\overline{\theta}^k}(k) = 0$.*

# 5 Proof of main result: Theorem 4.1

Consider in this section a sequence of evaluations $(\theta^k)_k$ that satisfies the LTC. As the proof is rather long, we divide it into two main parts:

- in Subsection 5.1, we present the first preliminary result, Proposition 5.1. It is used in two ways: first, we obtain an immediate consequence of it for later use, which bounds $\liminf_k V_{\theta^k}$ from below in terms of the auxiliary value functions $\{V_{\mathcal{T}_t\sharp\theta^k} : k \in \mathbb{N}^*, t \in \mathbb{R}_+\}$; second, we deduce from it in Corollary 5.2 the proof for Part (i) of Theorem 4.1.
- In Subsection 5.2, we prove Parts (ii)-(iii) of Theorem 4.1. Lemma 5.4 gives an upper bound of $\limsup_k V_{\theta^k}$ in terms of the auxiliary value functions $\{V_{\mathcal{T}_t\sharp\theta^k} : k \in \mathbb{N}^*, t \in \mathbb{R}_+\}$, which is, together with the result from Proposition 5.1, used to end the proof.

## 5.1 A first preliminary result and proof for Part (i)

**Proposition 5.1** *For any $\mu$ in $\Delta(\mathbb{R}_+)$, and any initial state $y_0$ in $\mathbb{R}^d$,*

$$\inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta}(y_0) \leq \liminf_k V_{\theta^k}(y_0).$$

*In particular, we have for all* $y_0$ *in* $\mathbb{R}^d$*,*

$$\sup_{k\in\mathbb{N}^*}\inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^k}(y_0) \le \liminf_k V_{\theta^k}(y_0).$$

**Proof:** Fixing $y_0$ and $\mu$, we set $\beta =_{def} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta}(y_0)$. For any $\varepsilon > 0$ fixed, there exists some $T_0 > 0$ such that $\mu([T_0,\infty)) < \varepsilon$. Take any control $u(\cdot)$ in $\mathcal{U}$. By definition of $\beta$, we have that

$$\forall T \ge 0,\ \int_{[0,+\infty)} g\left(y(t+T,u,y_0),u(t+T)\right)\mathrm{d}\mu(t) \ge \beta,$$

thus

$$\forall T \ge 0,\ \int_{[0,T_0]} g\left(y(t+T,u,y_0),u(t+T)\right)\mathrm{d}\mu(t) \ge \beta-\varepsilon. \tag{5.1}$$

For each $k \ge 1$, integrating both sides of (5.1) over $T\in[0,\infty)$ w.r.t. the evaluation $\theta^k$, we obtain

$$\int_{[0,+\infty)}\int_{[0,T_0]} g\left(y(t+T,u,y_0),u(t+T)\right)\mathrm{d}\mu(t)\mathrm{d}\theta^k(T) \ge \beta-\varepsilon. \tag{5.2}$$

Applying Fubini's Theorem to (5.2) yields

$$\beta-\varepsilon \le \int_{[0,T_0]}\left[\int_{[0,+\infty)} g\left(y(t+T,u,y_0),u(t+T)\right)\mathrm{d}\theta^k(T)\right]\mathrm{d}\mu(t) = \int_{[0,T_0]}\left[\gamma_{\mathcal{T}_t\sharp\theta^k}(y_0,u)\right]\mathrm{d}\mu(t), \tag{5.3}$$

where $\gamma_{\mathcal{T}_t\sharp\theta^k}(y_0,u) = \int_{[0,+\infty)} g\big(y(t+T,u,y_0),u(t+T)\big)\mathrm{d}\theta^k(T)$. According to Lemma 3.7, we have $|\gamma_{\theta^k}(y_0,u)-\gamma_{\mathcal{T}_t\sharp\theta^k}(y_0,u)| \le 2TV_t(\theta^k)$. This enables us to rewrite (5.3) as:

$$\begin{aligned}\beta-\varepsilon &\le \int_{[0,T_0]}\Big(\gamma_{\theta^k}(y_0,u)+2TV_t(\theta^k)\Big)\mathrm{d}\mu(t)\\ &\le \Big(\gamma_{\theta^k}(y_0,u)+2\overline{TV}_{T_0}(\theta^k)\Big)\mu([0,T_0])\\ &\le \gamma_{\theta^k}(y_0,u)+2\overline{TV}_{T_0}(\theta^k).\end{aligned}$$

The control $u(\cdot)\in\mathcal{U}$ being taken arbitrarily, we deduce that

$$\beta-\varepsilon \le V_{\theta^k}(y_0)+2\overline{TV}_{T_0}(\theta^k).$$

Since $(\theta^k)$ satisfies the LTC, $\overline{TV}_{T_0}(\theta^k)$ vanishes as $k$ tends to infinity. The proof is achieved. □

We end the proof for Part (i) of Theorem 4.1 by the following corollary of Proposition 5.1.

**Corollary 5.2** *[Proof for Part (i) of Theorem 4.1]*

$$\sup_{\mu\in\Delta(\mathbb{R}_+)}\inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\mu}(y_0) = \sup_{k\ge1}\inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\theta^k}(y_0),\ \forall y_0\in\mathbb{R}^d.$$

**Proof:** Fix $y_0 \in \mathbb{R}^d$, and denote $\varrho = \sup_{k\geq 1} \inf_{t\geq 0} V_{\mathcal{T}_t\sharp\theta^k}(y_0)$. It is clear that $\varrho \leq \sup_{\mu\in\Delta(\mathbb{R}_+)} \inf_{t\geq 0} V_{\mathcal{T}_t\sharp\mu}(y_0)$. Now for the converse inequality, consider for each $k \geq 1$, there exists $m(k)$ in $\mathbb{R}_+$ such that $V_{\mathcal{T}_{m(k)}\sharp\theta^k}(y_0) \leq \varrho + 1/k$. Since $\mathcal{T}_{m(k)}\sharp\theta^k$ – the image measure of $\theta^k$ by the function $s \mapsto s + m(k)$ – is also an evaluation on $\mathbb{R}_+$, we have:

$$\forall s \geq 0,\ TV_s(\mathcal{T}_{m(k)}\sharp\theta^k) = \sup_{Q\in\mathcal{B}(\mathbb{R}_+)} \Big|\theta^k\big((Q-m(k))\cap\mathbb{R}_+\big) - \theta^k\big((Q-m(k)+s)\cap\mathbb{R}_+\big)\Big| \leq TV_s(\theta^k) + \theta^k([0,s]).$$

We deduce that $(\mathcal{T}_{m(k)}\sharp\theta^k)_k$ satisfies the LTC whenever $(\theta^k)_k$ does so. According to Proposition 5.1, $\forall \mu \in \Delta(\mathbb{R}_+)$, $\inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\mu}(y_0) \leq \liminf_k V_{\mathcal{T}_{m(k)}\sharp\theta^k}(y_0) \leq \varrho$, thus $\sup_{\mu\in\Delta(\mathbb{R}_+)} \inf_{t\in\mathbb{R}_+} V_{\mathcal{T}_t\sharp\mu}(y_0) \leq \varrho$. The proof is complete. □

## 5.2 Proof for Parts (ii)-(iii)

In this subsection, we give the proof for Parts (ii)-(iii) of Theorem 4.1. We begin with the following result, which compares the values under evaluation $\mu$ and its $t$-"shifted" evaluation $\mathcal{T}_t\sharp\mu$ for any $t > 0$.

**Lemma 5.3** *Let $\mu$ in $\Delta(\mathbb{R}_+)$ be any evaluation. Then: for all $t \geq 0$ and $y_0 \in \mathbb{R}^d$,*

$$V_\mu(y_0) \leq V_{\mathcal{T}_t\sharp\mu}(y_0) + 2TV_t(\mu).$$

**Proof**: Fix $\mu \in \Delta(\mathbb{R}_+)$, $t \geq 0$, $y_0 \in \mathbb{R}^d$. By Lemma 3.7, we have

$$\gamma_\mu(y_0, u) \leq \gamma_{\mathcal{T}_t\sharp\mu}(y_0, u) + 2TV_t(\mu),\ \forall u(\cdot) \in \mathcal{U}.$$

For all $\varepsilon > 0$, take $u^\varepsilon(\cdot) \in \mathcal{U}$ be an $\varepsilon$-optimal control for $V_{\mathcal{T}_t\sharp\mu}(y_0)$, *i.e.*, $\gamma_{\mathcal{T}_t\sharp\mu}(y_0, u^\varepsilon) \leq V_{\mathcal{T}_t\sharp\mu}(y_0) + \varepsilon$. We obtain that

$$\gamma_\mu(y_0, u^\varepsilon) \leq V_{\mathcal{T}_t\sharp\mu}(y_0) + \varepsilon + 2TV_t(\mu).$$

Since $V_\mu(y_0) = \inf_{u(\cdot)\in\mathcal{U}} \gamma_\mu(y_0, u)$ and $\varepsilon > 0$ being arbitrary, we deduce that

$$V_\mu(y_0) \leq V_{\mathcal{T}_t\sharp\mu}(y_0) + 2TV_t(\mu),$$

which proves the lemma. □

The following result gives an upper bound on "$\limsup_k V_{\theta^k}$" in terms of the auxiliary value functions $\{V_{\mathcal{T}_t\sharp\theta^k} : k \in \mathbb{N}^*, t \in \mathbb{R}_+\}$.

**Lemma 5.4** *For all $T_0 \geq 0$ and any $y_0$ in $\mathbb{R}^d$,*

$$\limsup_k V_{\theta^k}(y_0) = \limsup_k \inf_{t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k}(y_0).$$

*In particular, for all $T_0 \geq 0$ and any $y_0$ in $\mathbb{R}^d$,*

$$\limsup_k V_{\theta^k}(y_0) \leq \sup_{k\in\mathbb{N}^*} \inf_{t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k}(y_0).$$

**Proof:** Fix $T_0 \geq 0$ and $y_0 \in \mathbb{R}^d$. The inequality $''\limsup_k \inf_{t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k} \leq \limsup_k V''_{\theta^k}$ is clear by taking $t = 0$ for each $k$. Now for the converse inequality $''\limsup_k \inf_{t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k} \geq \limsup_k V''_{\theta^k}$: according to Proposition 5.3, we have that for all $k$ and $t \leq T_0$,

$$V_{\theta^k}(y_0) \leq V_{\mathcal{T}_t\sharp\theta^k}(y_0) + 2TV_t(\theta^k).$$

For each $k \geq 1$, take $t^k \leq T_0$ with $V_{\mathcal{T}_{t^k}\sharp\theta^k}(y_0) \leq \inf_{0\leq t\leq T_0} V_{\mathcal{T}_{t^k}\sharp\theta^k} + \frac{1}{k}$, which gives us:

$$\begin{aligned} V_{\theta^k}(y_0) &\leq \inf_{0\leq t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k}(y_0) + \frac{1}{k} + 2TV_{t^k}(\theta^k) \\ &\leq \inf_{0\leq t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k}(y_0) + \frac{1}{k} + 2\overline{TV}_{T_0}(\theta^k). \end{aligned}$$

Since $(\theta^k)_k$ satisfies the LTC, $\overline{TV}_{T_0}(\theta^k)$ vanishes as $k$ tends to infinity. By taking "$\limsup_k$" on both sides of above inequality, the proof of the lemma is complete. □

Now we end the proof for Theorem 4.1. To do this, we first summarize results in Proposition 5.1 and Lemma 5.4 in the following chain form, which is then used for the study of the convergence of $(V_{\theta^k})_k$.

**Corollary 5.5** *For all $T_0 \geq 0$ and $y_0$ in $\mathbb{R}^d$,*

$$\sup_{k\geq 1}\inf_{t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k}(y_0) \geq \limsup_k V_{\theta^k}(y_0) \geq \liminf_k V_{\theta^k}(y_0) \geq \sup_{k\geq 1}\inf_{t\geq 0} V_{\mathcal{T}_t\sharp\theta^k}(y_0)$$

**Remark 5.6** *Corollary 5.5 states that the uniform convergence of $''\sup_{k\geq 1}\inf_{t\leq T_0} V_{\mathcal{T}_t\sharp\theta^k}''$ to $''\sup_{k\geq 1}\inf_{t\geq 0} V_{\mathcal{T}_t\sharp\theta^k}''$ as $T_0$ tends to infinity implies the uniform convergence of $(V_{\theta^k})_k$ as $k$ tends to infinity. Moreover, according to Corollary 5.2, in case of uniform convergence, the limit function is $V^*$.*

For any states $y$ and $\overline{y}$ in $\mathbb{R}^d$, let us define $\tilde{d}(y,\overline{y}) = \sup_{k\geq 1}|V_{\theta^k}(y) - V_{\theta^k}(\overline{y})|$. The space $(\mathbb{R}^d, \tilde{d})$ is now a *pseudometric* space (may not be Hausdorff).

The following is similar to the proof of Theorem 2.5 in [9], and is also similar to the proof of Theorem 3.10 in [10]. We rewrite it here for sake of completeness. Roughly speaking, we shall use the total boundedness of the space $\big(\{V_{\theta^k}\}, ||\cdot||_\infty\big)$ so as to deduce that the state space $(\mathbb{R}^d, \tilde{d})$ is totally bounded for the pseudometric metric $\tilde{d}$. This allows us to prove the convergence for $\tilde{d}$ of the reachable set $R^T$ to $R$ in bounded time. We are then able to prove the uniform convergence of "$\sup_{k\geq 1}\inf_{t\leq T_0} V_{\mathcal{T}_t\sharp\theta}$" to "$\sup_{k\geq 1}\inf_{t\geq 0} V_{\mathcal{T}_t\sharp\theta}$" as $T_0$ tends to infinity..

**Proof for Theorem 4.1, Parts (ii)-(iii)**.

We first prove Part (iii). One direction is clear: the uniform convergence of $(V_{\theta^k})$ implies the totally boundedness of the space $(\{V_{\theta^k}\}, ||\cdot||_\infty)$.

Let us prove the converse. Suppose that $(\{V_{\theta^k}\}, ||\cdot||_\infty)$ is totally bounded, so fixing any $\varepsilon > 0$, there exists a finite set of indices $I$ such that for all $k \geq 1$, there exists $i \in I$ satisfying

$$||V_{\theta^k} - V_{\theta^i}||_\infty \leq \varepsilon/3.$$

$\{\big(V_{\theta^i}(y)\big), y \in \mathbb{R}^d\}$ is a subset of the compact metric space $\big([0,1]^I, \|\cdot\|_\infty\big)$, thus it is itself totally bounded, so there exists a finite subset $X$ of $\mathbb{R}^d$ such that

$$\forall y \in \mathbb{R}^d, \exists x \in X, \forall i \in I, |V_{\theta^i}(y) - V_{\theta^i}(x)| \leq \varepsilon/3.$$

We have obtained that for each $\varepsilon > 0$, there exists a finite subset $X$ of $\mathbb{R}^d$ such that for every $y \in \mathbb{R}^d$, there is $x \in X$ satisfying: for any $k \geq 1$ there is some $i \in I$ with

$$\big|V_{\theta^k}(y) - V_{\theta^k}(x)\big| \leq \big|V_{\theta^k}(y) - V_{\theta^i}(y)\big| + \big|V_{\theta^i}(y) - V_{\theta^i}(x)\big| + \big|V_{\theta^i}(x) - V_{\theta^k}(x)\big| \leq \varepsilon,$$

thus $\tilde{d}(y,x) \leq \varepsilon$. This implies that the pseudometric space $(\mathbb{R}^d, \tilde{d})$ is itself totally bounded.

Fix now $y_0$ in $\mathbb{R}^d$. It is by definition that

$$\text{for all } \ T, S \in \mathbb{R}_+ \text{ with } S \geq T, \text{ we have } R^T(y_0) \subset R^S(y_0) \subset R(y_0),$$

and

$$\forall \bar{y} \in R(y_0), \ \exists \bar{T} > 0 \text{ with } \bar{y} \in R^{\bar{T}}(y_0).$$

From the totally boundedness of $(\mathbb{R}^d, \tilde{d})$, we show that $R^T$ converges to $R$ in the following sense

$$\forall \varepsilon > 0, \ \exists T \geq 0 : \ \forall \overline{y} \in R(y), \ \exists \tilde{y} \in R^T(y), \ \tilde{d}(\overline{y}, \tilde{y}) \leq \varepsilon. \tag{5.4}$$

Indeed, let us first take $\{y_\ell\}$ a finite $\varepsilon$-cover of $R(y_0)$ for $\tilde{d}$. For each $y_\ell$, put $T_\ell > 0$ with $y_\ell \in R^{T_\ell}(y_0)$. We then take $T = \max T_\ell$. Now for any $\bar{y} \in R(y_0)$, there is some $y_\ell$ with $\tilde{d}(\bar{y}, y_\ell) \leq \varepsilon$. Moreover, $y_\ell \in R^{T_\ell}(y_0) \subset R^T(y_0)$. This proves (5.4).

By Corollary 5.5, for all $T \geq 0$ $\big($ using $\inf_{\overline{y} \in R^T(y_0)} V_{\theta^k}(\overline{y}) = \inf_{t \leq T} V_{\mathcal{T}_t \sharp \theta^k}(y_0)$, cf. (2.4)$\big)$:

$$\sup_{k \geq 1} \inf_{\overline{y} \in R^T(y_0)} V_{\theta^k}(\overline{y}) \geq \limsup_k V_{\theta^k}(y_0) \geq \liminf_k V_{\theta^k}(y_0) \geq \sup_{k \geq 1} \inf_{\overline{y} \in R(y_0)} V_{\theta^k}(\overline{y}).$$

Consider $k \geq 1$ and $T \geq 0$ given by assertion (5.4) for the fixed $\varepsilon > 0$. Let $\overline{y} \in R(y_0)$ be such that $V_{\theta^k}(\overline{y}_0) \leq \inf_{\overline{y} \in R(y)} V_{\theta^k}(\overline{y}) + \varepsilon$, and then $\tilde{y}$ in $R^T(y_0)$ be such that $\tilde{d}(\overline{y}, \tilde{y}) \leq \varepsilon$. Since $V_{\theta^k}$ is clearly 1-Lipschitz for $\tilde{d}$, we obtain $V_{\theta^k}(\tilde{y}) \leq \inf_{\overline{y} \in R(y_0)} V_{\theta^k}(\overline{y}) + 2\varepsilon$. Consequently, $\inf_{\overline{y} \in R^T(y_0)} V_{\theta^k}(\overline{y}) \leq \inf_{\overline{y} \in R(y_0)} V_{\theta^k}(\overline{y}) + 2\varepsilon$ for all $k$, so

$$\sup_{k \geq 1} \inf_{\overline{y} \in R^T(y_0)} V_{\theta^k}(\overline{y}) \leq \sup_{k \geq 1} \inf_{\overline{y} \in R(y_0)} V_{\theta^k}(\overline{y}) + 2\varepsilon.$$

One obtains that $\limsup_{k \geq 1} V_{\theta^k}(y_0) \leq \liminf_{k \geq 1} V_{\theta^k}(y_0) + 2\varepsilon$, and so $\big(V_{\theta^k}(y_0)\big)_k$ converges. Since $(\mathbb{R}^d, \tilde{d})$ is totally bounded and all $V_{\theta^k}$ are 1-Liptschitz, the convergence is uniform.

Next, Part (ii) can be deduced from the proof of Part (iii). Let $(\theta^{\varphi(k)})$ be any subsequence of $(\theta^k)$ that converges uniformly to some function $V$. This implies that $(\{V_{\theta^{\varphi(k)}}\}, || \cdot ||_\infty)$ is totally bounded. As we have shown in the proof of Part (iii) that if $(\{V_{\theta^{\varphi(k)}}\}, || \cdot ||_\infty)$ is totally bounded, $(V_{\theta^{\varphi(k)}})$ converges uniformly to $V = V^*$, which implies Part (ii) that $V^*$ is the unique accumulation point (for the uniform convergence) of the sequence $(V_{\theta^k})_k$. $\square$

# 6 Discussion on a weaker long-term condition

Below is a weaker form of the long-term condition (LTC):

**Long-term condition' (LTC')** A sequence of evaluations $(\theta^k)_{k\geq 1}$ satisfies the LTC' if:

$$\forall s > 0, \quad TV_s(\theta^k) \xrightarrow[k\to\infty]{} 0. \tag{6.1}$$

It is unclear whether the LTC' is strictly weaker than the LTC or not. One might want to construct an example of $(\theta^k)_k$ such that $TV_s(\theta^k) \xrightarrow[k\to\infty]{} 0$ for all $s > 0$ while $\overline{TV}_{s_0}(\theta^k) \xrightarrow[k\to\infty]{} \alpha > 0$ for some $s_0 > 0$ and $\alpha > 0$. The following example shows that this is possible if we consider only $s$ being rational numbers. In general, the question is still open.

**Example 6.1** *Given a positive integer $k$, consider the density $\theta^k$ with support included in $[0, k]$ by dividing $[0, k]$ in $k^2$ consecutive small intervals of length $1/k$, and $\theta^k$ is uniform over the union of all small odd intervals... and puts no weight on even small intervals. Define the support*

$$S_k = \bigcup_{l\in\mathbb{N}, l\leq \frac{k^2-1}{2},} \left[\frac{2l}{k}, \frac{2l+1}{k}\right).$$

*$\theta^k$ has density:*

$$f_k(x) = \frac{2}{k}\,\mathbb{1}_{x\in S_k} = \frac{2}{k}\,\mathbb{1}_{x\in[0,k], E(kx)\in 2\mathbb{N}}$$

*(where $2\mathbb{N}$ is the set of even numbers in $\mathbb{N}$, $E(x)$ is the integer part of $x$).*

*For each $k$, we have (consider $s = 1/k$):*

$$\sup_{0\leq s\leq 1} \int_{x\geq 0} \big|f_k(x+s) - f_k(x)\big| \mathrm{d}x \geq 2 - 1/k$$

*Consider now only $k$ of the form $n!$, and we define the density $g_n = f_{n!}$ for each $n$ in $\mathbb{N}$. For all $x \geq 0$,*

$$g_n(x+s) - g_n(x) = \frac{2}{n!}\left(\mathbb{1}_{E(n!(x+s))\in 2\mathbb{N}, x+s\leq n!} - \mathbb{1}_{E(n!x)\in 2\mathbb{N}, x\leq n!}\right).$$

*Assume $s$ is a rational number. Then for $n$ large enough, $n!s$ is an even integer, so for all $x$ such that $0 \leq x \leq n! - s$, we have $g_n(x+s) - g_n(x) = 0$. Consequently,*

$$\int_{x\geq 0} \big|g_n(x+s) - g_n(x)\big| \mathrm{d}x \xrightarrow[n\to\infty]{} 0.$$

# 7 Appendix

**Proof for Lemma 1**: The following computation of $f'_\theta(t)$ is straightforward:

$$\forall t > 0, \quad f'_\theta(t) = \frac{1}{\sigma\sqrt{2\pi}}\Big[\exp\left(-\frac{1}{2}\Big(\frac{t-m}{\sigma}\Big)^2\right)\frac{m-t}{\sigma^2} - \exp\left(-\frac{1}{2}\Big(\frac{t+m}{\sigma}\Big)^2\right)\frac{m+t}{\sigma^2}\Big],$$

thus

$$f'_\theta(t) > 0 \ (resp. < 0) \iff (m-t)\exp\left(-\frac{1}{2}\Big(\frac{t-m}{\sigma}\Big)^2\right) - (m+t)\exp\left(-\frac{1}{2}\Big(\frac{t+m}{\sigma}\Big)^2\right) > 0 \ (resp. < 0).$$

As a consequence, one obtains that $f'_\theta(t) < 0, \ \forall t \geq m$. Now we look at $t \in (0, m)$. Denote $H(t) =_{def} \exp\left(\frac{2mt}{\sigma^2}\right) - \frac{m+t}{m-t}$, which enables us to write:

$$f'_\theta(t) > 0 \ (resp. < 0) \iff H(t) > 0 \ (resp. < 0), \quad \forall t \in (0, m).$$

From the above analysis, we deduce that the proof of the lemma is reduced to the proof for

**Claim** *There is some $t^* \in [0, m)$ such that $H(t) < 0$ for $t \in (0, t^*)$ and $H(t) > 0$ for $t \in (t^*, m)$. Moreover, such $t^*$ satisfies $(t^*)^2 \geq m^2 - \sigma^2$.*

In order to prove the claim, we compute

- the values at the end point: $H(0) = 0$ and $\lim_{t \to m^-} H(t) = -\infty$;
- the first-order derivative at any $t \in [0, m)$:

$$H'(t) = \exp\left(\frac{2mt}{\sigma^2}\right)\frac{2m}{\sigma^2} - \frac{2m}{(m-t)^2} \tag{7.1}$$

- at any rest point $t^e \in [0, m)$ (*i.e.*, $H(t^e) = 0$):

$$\exp\left(\frac{2mt^e}{\sigma^2}\right) = \frac{m+t^e}{m-t^e},$$

  which is substituted back into (7.1), to yield

$$H'(t^e) > 0 \ \big( \ resp. \ H'(t^e) < 0 \ \big) \iff (t^e)^2 < m^2 - \sigma^2 \ \big( \ resp. \ (t^e)^2 > m^2 - \sigma^2 \ \big). \tag{7.2}$$

Next, it is easy for us to prove the following result:

*Let $t_1^e \in [0, m)$ be a rest point for $H(\cdot)$, and suppose that $t_2^e \in (t_1^e, m)$ is the smallest rest point after $t_1^e$. Then $H'(t_1^e)H'(t_2^e) \leq 0$ and if $H'(t_1^e) \leq 0$, such $t_2^e$ does not exist.*

Indeed, $H'(t_1^e)H'(t_2^e) \leq 0$ can be derived from the continuity of $H(\cdot)$; suppose that $H'(t_1^e) \leq 0$, we have from (7.2) that $(t_1^e)^2 \geq m^2 - \sigma^2$ and $H'(t_2^e) \geq 0$, thus $(t_2^e)^2 \leq m^2 - \sigma^2$. However, this leads to a controdiction to $t_2^e > t_1^e$, so $t_2^e$ does not exist whenever $H'(t_1^e) \leq 0$.

Finally, remark that $H(0) = 0$, thus $t = 0$ is a rest point. We discuss the following two cases:

Case 1. $m^2 - \sigma^2 \leq 0$, thus $H'(0) \leq 0$.

This implies that no rest point exists after 0. Since $\lim_{t \to m^-} H(t) = -\infty$, we deduce that $H(t) < 0, \ \forall t \in (0, m)$. The claim is proved for $t^* = 0$.

Case 2. $m^2 - \sigma^2 > 0$, thus $H'(0) > 0$.

$\lim_{t \to m^-} H(t) = -\infty$ implies that some rest point exists in $(0, m)$. Take $t^e$ the closest to 0, implying that $H(t) > 0, \ \forall t \in (0, t^e)$. Further, we obtain that $H'(t^e) \leq 0$ by the continuity of

$H(\cdot)$. Again, there exists no other rest point after $t^e$. Since $\lim_{t\to m^-} H(t) = -\infty$, we deduce that $H(t) < 0$, $\forall t \in (t^e, m)$. The claim is proved for $t^* = t^e$.

To conclude, we see that in both cases such $t^*$ exists and satisfies $(t^*)^2 \geq m^2 - \sigma^2$, thus the claim is proved. This finishes our proof for the lemma. □

# References

[1] O. Alvarez and M. Bardi. Ergodicity, stabilization, and singular pertubations for Bellman-Isaacs equations. *Mem. Amer. Math. Soc.*, **204**(2010), no. 960.

[2] M. Arisawa. Ergodic problem for the Hamilton-Jacobi-Belmann equations. *Ann. Henri Poincaré, Analyse Nonlinéaire*, **15**(1998), 1–24.

[3] M. Arisawa and P.L. Lions. On ergodic stochastic control. *Comm. Partial Differential Equations*, **23**(1998), 2187–2217.

[4] A. Bensoussan. *Perturbation Methods in Optimal Control.* Wiley/Gauthiers-Villas, Chichester, 1988.

[5] L. Devroye and L. Györfi. *Nonparametric Density Estimation: the L1 view.* Wiley, New York, 1985.

[6] V. Gaitsgory. On the use of the averaging method in control problems. *(Russian) Differentsialnye Uravneniya*, **22** (1986), 1876–1886.

[7] R.Z. Khasminskii. On the averaging principle for Itô stochastic equations. *Kybernetika*, **4**(1968), 260–279.

[8] M. Quincampoix and J. Renault. On the existence of a limit value in some nonexpansive optimal control problems. *SIAM Journal on Control and Optimization*, **49**(2011), 2118-2132.

[9] J. Renault. General long-term values in dynamic programming. *Journal of Dynamics and Games*, **1**(2014), 471–484.

[10] J. Renault. Uniform value in dynamic programming. *J. Eur. Math. Soc.(JEMS)*, **13**(2011), 309–330.

[11] J. Renault and X. Venel. A distance for probability spaces, and long-term values in markov decision processes and repeated games. *arXiv:1202.6259*, 2013.

[12] S. Sorin. *A First Course on Zero-sum Repeated Games.* Springer, 2002.

[13] B. Ziliotto. General limit value in stochastic games. *arXiv:1410.5231*, 2014.

*Email address*: xxleewhu@gmail.com
*Email address*: Marc.Quincampoix@univ-brest.fr
*Email address*: jerome.renault@tse-fr.eu